





Quick Guide



Agilia® SP MC WiFi Connected Generation NOTE: This Quick Reference Guide is not intended to be comprehensive instructions for the set-up and operation of the Agilia SP MC and Agilia SP MC WiFi Volumetric Infusion Pumps.






INSTALLING A SYRINGE

Connect the extension set to the syringe according to local practices. Prime and check that there is no air in the syringe or extension set.

IMPORTANT: Ensure that the patient is not connected.

- Place the syringe into the cradle, the flanges inserted in the provided slot.
- Secure the syringe with the syringe barrel clasp.
- Move the plunger driver gently until it makes contact with the plunger head.



General Operations

1. Selecting a Profile



- Press <ON/OFF> to power ON the pump.
- Use arrow keys to select desired profile.
- Press <OK> to confirm the profile.

2. Selecting a Syringe



- Press <OK> to confirm the displayed syringe, or <C> to access the syringe list.
- Press the arrow keys to select a new syringe.



Press <OK> to confirm the new syringe.

3. Selecting a Drug



Press the arrow keys to scroll to the drug's first letter, and press <OK>.

NOTE: Drugs are sorted alphabetically by the first letter.



- Press the arrow keys to scroll to the drug name, and press < OK>.
- Press <OK> to acknowledge the clinical advisory message and continue programming, or <C> to change the drug.

Programming an Infusion by Flow Rate



- Press the arrow keys to program the flow rate.
- Press <start>.

Programming an Infusion by Dose

Selecting the Drug Concentration





Finite values

- Press the arrow keys to select the concentration.
- Press <OK> to confirm.

Selecting the Patient's Characteristics



- Press the arrow keys to enter the patient's weight or body surface area.
- Press <OK> to confirm.

Programming the Infusion



- Press the arrow keys to program the dose rate value.
- Press <OK> to confirm.

Starting an Infusion

Connect the administration set to the patient's IV access device using aseptic technique and according to your healthcare facility policy.





v rate Dose

■ Press <start> to start the infusion, or <C> to modify the infusion settings.

Functions During Infusion

1. Stop infusion



- Press <STOP> key: After 2 minutes, an alarm is generated as a reminder that the infusion is stopped.
- Confirm or modify settings prior to restarting the infusion.

2. Rate Titration (while infusing)



CARBoplatin 2 mg/ml.

Altia

Billia

B

- Press the arrow keys to modify the flow rate or dose.
- Press <OK> to confirm.

Accessing the Menu Functions

- Press <MENU> key to access menu items, listed on left side of display.
- Use arrow keys to select menu item.
- Press <enter> to activate menu function.
- Follow prompts on the screen.

Powering off



- Press <STOP> key to stop the infusion.
- Press and hold <ON/OFF> until the pump powers off.

Administering a Bolus: Direct or Programmed Bolus

There are two ways to deliver a bolus during an infusion:

1. Direct Bolus



- During the infusion, press <BOLUS> key.
- Press <bol>
 Press <bol>
 bolus> to confirm access.



- Press and hold <BOLUS> key.
- To stop the bolus, release the <BOLUS> key: The infusion resumes its previous rate.

2. Programmed Bolus

- Press <BOLUS> key.
- Press <prog>.



Flow rate



Dose

- Press the arrow keys to program the bolus volume or dose, and press <OK>.
- Press the arrow keys to program the bolus duration, and press <OK>: The flow rate is calculated automatically.



Flow rate



Dose

■ Press <start> to start the bolus.

After bolus is delivered the infusion resumes its previous rate.

Interrupting a Programmed Bolus



Flow rate



■ Press <STOP> key.

- Press <no> to return to the infusion.
- Press <start> to continue the bolus.



KEYPAD DESCRIPTION On / Off Increment)(fast/low) Decrement (low/fast) Confirm Value / Move to Next Field Stop / Pause Bolus / Prime (Pressure Menu/ Menu Cancel Value / Move Back to Previous Field Alarm Silence LIGHT INDICATORS **-**(3) **Power Supply Indicator Battery Charge Status Indicator** Infusion in Progress: flashing green Low-Priority Alarm: constant yellow / Medium-Priority Alarm: flashing yellow **********

High-Priority Alarm: flashing red

This document may contain inaccuracies or typographical errors. Modifications may thus be made, and included in later editions. Due to the evolution of standards, and of legal texts and materials, the characteristics indicated in the text and images of this document are applicable only to the device with which it is included.

